

## Selection criteria

**Selection period:** 2015-2024

**Database extracted** in Oct-2025

*Data collected from patient medical files  
Quality control processes and yearly updates*

**69 326** patients in the database

## Median follow-up

*Calculated from advanced or metastatic diagnosis, according to reverse Kaplan-Meier method*

**60.5** months

## Minimum estimated follow-up

24 months for **50 926** pts

36 months for **44 553** pts

**7 664** patients with **SCLC**

**56 470** patients with **NSCLC**

**50 982**

patients with **advanced or metastatic NSCLC**

**42 196**


patients with advanced or metastatic NSCLC who received a **first line of treatment**

**11 488**


patients with **unresectable stage III NSCLC**

- Female or male patients
- $\geq 18$  years old
- who received a first treatment (systemic or radiotherapy) for lung cancer or diagnosed with metastatic lung cancer at one of the 38 participating French medical centers

**♂ 44 955 ♀ 24 371**




**Median age at initial diagnosis**  
**66** years



**61 283** patients with **advanced or metastatic** lung cancer

**21 599** patients with **oncogenic drivers**



**PD-L1** screening performed in **37 387** pts

**TPS  $\geq 1\%$**  in **23 151** patients  
**TPS  $\geq 50\%$**  in **11 141** patients